



Hand Hygiene Policy

Version 12: September 2022

First Issued July 2004

Review date September 2024

Contents

(For quick access to a specific heading - press CTRL and click your mouse to follow the link for the below options)

1.	INTRODUCTION	Error! Bookmark not defined.
2.	PURPOSE AND SCOPE.....	Error! Bookmark not defined.
3.	RESPONSIBILITIES	Error! Bookmark not defined.
4.	POLICY STATEMENT	3
5.	PROCEDURE	3
6.	MONITORING AND REVIEW	6
7.	REFERENCES	7
8.	AUTHOR	8
9.	APPENDICES	8
	Appendix 1 Hand Hygiene Technique	9
	Appendix 2 Hand Hygiene Audit Tool	10
	Appendix 3 Hand Hygiene Disciplinary Framework	11
	Appendix 4 5 Moments of Hand Hygiene at the Point of Care	12
	Appendix 5 5 Moments of Hand Hygiene at the Point of Care part 2	13
10	EQUALITY AND DIVERSITY IMPACT ASSESSMENT	14
11.	DOCUMENT CONTROL	14

1. INTRODUCTION

Hand hygiene is one important component in the battle against cross-infection. Minimising risks of infection to patients depends on a range of factors. However, just by increasing hand hygiene alone can dramatically reduce the risk of a patient acquiring an infection. Scientific evidence demonstrates that the bacteria that causes healthcare associated infections (HCAI) are most frequently spread from one patient to another on the hands of healthcare workers. Health Act 2006 (2008), The Health and Social Care Act 2008 (2015), Essential Steps to Safe, Clean Care (2007), Cleanyourhands (2008) Epic 3 (2014). Therefore, hand hygiene carried out immediately before and after contact with patients i.e. at point of care, is the best way to prevent HCAs.

Hand hygiene procedures do not only relate to clinical staff, all staff must wash their hands correctly and at the appropriate time, see WHO 'your 5 moments for hand hygiene'.

2. PURPOSE AND SCOPE

To ensure correct hand washing practice. This document applies to all staff either employed or contracted by East Coast Community Healthcare (ECCH). These staff may work within our premises, patients own homes, or care settings owned by other agencies.

3. RESPONSIBILITIES

It is the responsibility of all staff to ensure that they adhere to evidence based best practice. All staff must take responsibility for their own hand decontamination and should act as an advocate for all their clients and others to ensure that everyone decontaminates their hands appropriately.

4. POLICY STATEMENT

This guidance will be implemented to ensure adherence to safe practice

5. PROCEDURE

Thorough hand washing is undoubtedly one of the simplest and most effective ways of preventing person-to-person transmission of infective agents in clinical practice.

Hand decontamination has a dual role in protecting both the patient and the healthcare worker. Hands readily pick up and transfer micro-organisms. Hands must be decontaminated between all activities that result in even superficial contact with patient surroundings.

There is no set frequency for hand decontamination as it is determined by clinical actions, but there are guidelines from the World Health Organisation (WHO) and the National Patient Safety Agency on 'your 5 moments for hand hygiene at the point of care' (see 14 & 15). A risk assessment of the activity intended to be performed, will determine the appropriate decontamination process and the choice of cleansing agent.

There should be an appropriate use of 'single use non sterile gloves' and 'sterile gloves' within ECCH. Hand hygiene should be performed before and after the use of gloves. Gloves should not be cleaned with hand sanitizer/rub they must be changed between each patient contact or episode of care.

The principles of hand decontamination apply equally to healthcare provided in hospitals as they do to care provided in the community, but these may need to be adapted to suit local circumstances.

The Microbiology of the hand

There are two populations of microbes present on the hands.

Transient micro-organisms – these superficially present on the skin and are:

- Easily removed by routine hand washing
- Easily acquired by touch
- Readily transferred to the next person or surface touched
- The usual source of cross infection

Resident micro-organisms – these are deep seated within the skin:

- They are difficult to remove
- Are not readily transferred to other people or surfaces
- Can enter tissues and establish infection during highly invasive procedures such as surgery
- Play an important part in protecting skin from other harmful organisms

Routine hand decontamination

The aim of this is to remove transient micro-organisms before they can be transferred. Hands that are visibly soiled with dirt or organic material, or potentially contaminated with micro-organisms should be washed using liquid soap and water using the seven step technique taking 15-30 seconds, then rinsed and dried thoroughly. Antibacterial hand cleansing solutions/soap are not routinely recommended as they kill off the resident micro-organisms, (the ones which help protect the skin), as well as the transient ones.

If hands are potentially contaminated, but visibly clean they may be decontaminated at point of care, using an alcohol based preparation, and the seven step technique taking 15-30 seconds, until both hands are dry. There are two makes of hand sanitizer which are approved for use within ECCH, they are Gojo Purell or Ecolab Spirigel.

Requirements

Estates and Facilities.

Sinks specifically designated to facilitate effective hand decontamination should be provided in all clinical areas, with elbow or non-touch taps and which conform to current recommendations, a supply of warm water, liquid soap and disposable paper towels in wall mounted dispensers, and a foot operated, closed lidded disposal bin for household waste. It is preferable to use products available via the approved supply chain when purchasing soap/paper towels/hand cream/hand rub/hand gel.

Hand sanitizer should be available at the point of care, preferably wall or bed/locker mounted at strategic points within clinical areas e.g. on notes trolley for during ward rounds. It is the responsibility of members of staff who finish the end of a bottle/container to replenish with fresh stock. Hotel Services will be responsible for daily cleaning, maintenance and reporting faulty hand hygiene products/stations. It is the responsibility of the ward to keep check list of cleaning and maintenance.

Hand sanitizer does not replace the need for conveniently located and dedicated facilities for hand washing in clinical areas, and where possible extra sinks will be fitted which conform to current recommendations.

Single use patient hand wipes must be available for those patients who are unable to access liquid soap and water for hand washing e.g. before meals, after using the toilet.

There should be hand sanitizer/bottles of liquid soap/hand wipes carried as individual dispensers for specialist areas or special circumstances e.g. community, paediatrics, domiciliary visits. Dispensers must not be refilled.

In areas where it is considered unsafe to have hand sanitizer easily available, a risk assessment must be carried out and submitted.

At the entrance to all in patient healthcare facilities there should be signs explaining the importance of hand hygiene and the ECCH's commitment to improving hand hygiene. All hand hygiene stations whether sinks or hand sanitiser, should have easily visible, clear signage which should encourage staff, patients and visitors to comply with hand hygiene measures.

Preparation of hands:

Intact skin is an effective barrier to prevent micro-organisms entering the body. Therefore, all cuts, abrasions and other skin lesions on the hands (and other exposed areas of skin) of health care workers should be covered with an occlusive waterproof dressing.

Best practice in hand hygiene consists of keeping fingernails short and clean, (when holding hands up, palm facing you, you should not be able to see the white of the nails). Best practice requires removal of all jewellery, except one plain ring band (please also refer to the Uniform Policy, and Bare Below the Elbows in Clean, Safe Care. DH (2008) Epic 3 (2014)) including nail jewellery, nail polish and artificial fingernails.

Hand creams should be regularly applied to hands to protect the skin from the drying effects of regular hand decontamination. Communal jars of hand cream are not advisable as the contents may themselves become contaminated and therefore become a source of cross-infection. Wall mounted pump dispensers and hand cream are available through stores.

Nail brushes are not recommended for routine use as they can damage the skin. Where nail brushes are used they should be sterile and single use only.

When to decontaminate hands:

Hands should always be cleaned:

- Before starting and at the end of each work period.
- Before and after each 'hands on' patient contact at point of care.
- Before and after carrying out each aseptic procedure.
- After any contact with body fluids or secretions.
- After handling soiled or contaminated equipment or linen.
- Before and after administering drugs.
- Whenever skin is visibly soiled.
- Before and after use of gloves.
- Before performing or assisting at operative procedures, a surgical scrub for hand decontamination should be performed.
- After using the lavatory.
- Before eating, drinking or handling food.
- After contact with patient surroundings.

This list is not exhaustive and we expect all staff to use the 5 moments for hand hygiene charts and their clinical judgement to decide appropriateness.

Cleansing agents

- Plain liquid soap and water is sufficient for most routine daily activities. The seven step procedure for cleaning hands should be used. Hand washing with liquid soap and water suspends the micro-organisms in solution and allows them to be rinsed off – this is referred to as *mechanical removal* of micro-organisms. Liquid soap should be used, with enough soap applied to ensure that hands are well lathered all over. The dispenser should be wall mounted and regularly maintained, with individual replacement cartridges that are discarded when empty. There should be nominated staff to be responsible for this.
- Alcohol-based hand rub should be used to decontaminate hands before and after direct patient contact and clinical care, except in the following situations when soap and water must be used:

When hands are visibly soiled or potentially contaminated with body fluids and/or when caring for patients with vomiting or diarrhoeal illness, regardless of whether or not gloves have been worn. (Loveday et al. 2014)

- It should be applied using an evidence-based technique, we recommend the seven step procedure for cleaning hands, and about a 3ml dose dispensed (1 metered dose of a 800 or 1000ml pump container) should be used until both hands are dry. **Hand sanitiser should not be used when there is diarrhoea or vomiting as it is less effective against some organisms than washing with liquid soap and water.** Hand sanitiser can be used consecutively until the hands start to feel tacky when they should be washed with liquid soap and water.
- Staff experiencing problems with skin irritation or with concerns should contact the relevant Occupational Health Department – 01502 445300

Surgical hand decontamination

The aim of this is to substantially reduce resident micro-organisms and remove or destroy transient micro-organisms.

Used prior to surgical or other highly invasive procedures where extra care must be taken to prevent micro-organisms on hands being introduced into tissues should gloves become damaged.

This process is achieved by using an antiseptic hand cleansing preparation.

- Antiseptic hand washing solutions used with water will remove and destroy micro-organisms by the *chemical removal* of micro-organisms.
- Hand disinfection carried out in this way will reduce counts of colonising resident flora as well as removing or destroying transient micro-organisms. Some have a residual activity providing continued anti-microbial activity, which is of benefit during surgical procedures. Examples of aqueous antiseptic solutions are: chlorhexidine, iodophors and triclosan.

Patients and Visitors

Staff should educate and encourage patients to decontaminate their hands after toileting, before consumption of food or drink and before caring for invasive lines or dressings.

Visitors should be encouraged to decontaminate their hands when entering or leaving wards/departments and before and after contact with patients.

6. MONITORING AND REVIEW

It is the responsibility of all department heads/professional leads to ensure that staff they manage adhere to this guidance. All managers are required to ensure their teams complete the audit.

The target for correct and complete hand hygiene is set at 100% and should be monitored through monthly ward audits and quarterly departmental audits.

Should a ward area fall below 80% on 2 consecutive months the ward manager and the Infection Prevention & Control link nurse from the area will, using reflection, produce a written action plan using SMART objectives, (S-specific, M-measurable, A-achievable, R-realistic and T-timely). This will be submitted to Matrons and to the Infection Prevention & Control Team (IP&C) and will be taken to the Infection Prevention & Control Committee (IPACC).

The IP&C Team will set the number of hand hygiene audits to be carried out in each area depending on staff numbers and whole time equivalents. The audit should be carried out using the appropriate form found in this document. This should be returned monthly to the IP&C Team. These results are reported to IPACC on a quarterly basis. They are included in the Infection Control Quarterly and Annual reports which are submitted to the board and provide part of the assurance required by ECCH to the Care Quality Commission of adherence to the Health and Social Care Act 2008 (2015) (DH 2009).

The IP&C Team will carry out induction and yearly mandatory infection control training for clinical staff and 3 yearly for non-clinical staff. It is the responsibility of all ECCH staff to maintain their compliance with Infection Prevention and Control training, in person or on-line. The IP&C Team will also observe hand hygiene technique and opportunities during regular community hospital visits and during audits for all ECCH areas.

This guidance will be reviewed every two years by the Infection Prevention and Control Team, unless a substantive change occurs before this date.

7. REFERENCES

- Ayliffe GAC, Fraise AP, Geddes AM, Mitchell K [2000] *Control of Hospital Infection- a practical handbook*. 4th edition. Arnold.
- Department of Health (2007) *Essential Steps to Safe clean Care*. DH Publications. London.
- Department of Health. (2007) *Bare Below the Elbows*. Johnson outlines new measure to tackle hospital bugs. Johnson A London.
- Department of Health (2008) *Clean, safe care: reducing infections and saving lives*. DH Publications. London.
- Department of Health (2008) *The Health Act 2006: Code of Practice for the prevention and Control of Healthcare Associated Infections*. DH Publications. London.
- Department of Health (2015) *The Health and Social Care Act 2008 : Code of Practice for the NHS on prevention and control of healthcare associated infections and related guidance*. London
- Department of Health (2010) *Uniforms and Workwear: Guidance on Uniform and Workwear Policies for NHS Employers*. DH London
- Hand washing liaison group [1999]. *A modest measure with big effects*, *British Medical Journal* 318:686
- ICNA [2002] *Hand Decontamination Guidelines*
- Jacobson G et al [1985]. *Handwashing: ring wearing and number of micro-organisms*.
- Loveday, HP (2014) *Epic 3: National Evidence-Based Guidelines for Preventing Healthcare-Associated Infections in NHS Hospitals in England*. J.A Wilson, R.J Pratt, M. Golsorkhi, A Tingle, A Bak, J. Browne, J. Prieto, M.Wilcox. *Journal of Hospital Infection* 86S1 (20914) S1-S70. On-line at www.sciencedirect.com
- National Institute for Health and Clinical Excellence [2017] *Infection prevention and control of healthcare-associated infections in primary and community care*. NICE clinical guideline 139 www.nice.org.uk/cg139 (accessed 21/03/2022)

National Institute for Health and Clinical Excellence [2014] Infection prevention and control..NICE quality standard QS61 April 2014 www.nice.org.uk/qs61
National Patient Safety Agency [2008] Cleanyourhands Campaign
Reybrouk G [1983]. Role of hands in the spread of nosocomial infections. Journal of Hospital Infection 4:103-110
RCN [2017] Essential Practice for Infection Prevention and Control
World Health Organisation (2009) WHO Guidelines on Hand Hygiene in Health Care: First Global Patient Safety Challenge. Clean Care is Safer Care. World Health Organisation, Geneva.
World Health Organisation (2012) Hand Hygiene in Outpatient and Home-based care and Long-term care facilities. World Health Organisation. Geneva.
Whqlibdoc.who.int/publications/2009/9789241597906_eng.pdf

8. AUTHOR

Infection Prevention and Control Team, East Coast Community Healthcare

9. APPENDICIES

Appendix 1

Research has shown that the technique is as important as the time taken and the agent used

Hand Hygiene Technique



Rub hands palm to palm



Rub back of each hand with the palm of the other



Rub palm to palm with fingers interlaced



Rub with back of fingers to opposing palms with fingers interlocked



Rub tips of fingers in opposite palm in circular motion



Rub each thumb clasped in opposite hand using rotational movement



Rub each wrist with opposite hand

Each step should be repeated 5 times.


This same seven step technique should be used when using soap and water or hand sanitiser.

Infection Prevention and Control Team
East Coast Community Healthcare
Tel: 01502 445361
Email: infectionprevention@ecchcic.nhs.uk

Appendix 2

Hand Hygiene Audit Tool mk 5				Site;		Date;												
Obs	Designation of staff	Description of Task (Why you are washing your hands. dressings, feeding, personal care)	Hand Hygiene please state opportunity	7 Step hand hygiene technique										Hand Care			Personal Protective Equipment	Signature
				wetting hands/correct amount of gel	palm to palm	palm to back of hands	palm to palm fingers interlaced	back of fingers to palm	thumbs	rotational rubbing fingertips	wrists	rinsing	drying	Minimal Jewellery (1 plain band/no wrist watch)	Skin Condition hands (intact/cuts covered)	Finger nails clean, short, no varnish		
			See Key	Yes/No	Yes/No	Yes/No	Yes/No	Yes/No	Yes/No	Yes/No	Yes/No	Yes/No	Yes/No	Yes/No	Yes/No	Yes/No	Yes/No/N/A	
1																		
2																		
3																		
4																		
5																		
6																		
7																		
8																		
9																		
10																		
11																		
12																		
Total																		
% Compliance																		
Return to Research & Audit Team, ECCH HQ by end of the month												Key	1 - Before contact					
													2 - After contact					
Designation of staff - N-Nurse, Dr- Doctor, P-Physio, OT-OT, B/A- Bank/Agency													3 - Before Aseptic Technique					
Adapted from NHS Leeds ESSOC Hand Hygiene Audit Tool													4 After Body Fluid Exposure Risk					
Infection Prevention Team													5 - After contact with patient Surroundings					
ECCH 24/04/2012													6 - Missed opportunity					

Hand Hygiene Disciplinary Framework




Your 5 moments for hand hygiene at the point of care



1	BEFORE PATIENT CONTACT	WHEN? Clean your hands before touching a patient when approaching him/her WHY? To protect the patient against harmful germs carried on your hands
2	BEFORE AN ASEPTIC TASK	WHEN? Clean your hands immediately before any aseptic task WHY? To protect the patient against harmful germs, including the patient's own, from entering his/her body
3	AFTER BODY FLUID EXPOSURE RISK	WHEN? Clean your hands immediately after an exposure risk to body fluids (and after glove removal) WHY? To protect yourself and the healthcare environment from harmful patient germs
4	AFTER PATIENT CONTACT	WHEN? Clean your hands after touching a patient and her/his immediate surroundings when leaving the patient's side WHY? To protect yourself and the healthcare environment from harmful patient germs
5	AFTER CONTACT WITH PATIENT SURROUNDINGS	WHEN? Clean your hands after touching any object or furniture in the patient's immediate surroundings when leaving - even if the patient has not been touched WHY? To protect yourself and the healthcare environment from harmful patient germs

Adapted from WHO World Alliance for Patient Safety 2006

Your 5 moments for hand hygiene at the point of care



1	BEFORE PATIENT CONTACT	WHEN? Clean your hands before touching a patient when approaching him/her WHY? To protect the patient against harmful germs carried on your hands
2	BEFORE AN ASEPTIC TASK	WHEN? Clean your hands immediately before any aseptic task WHY? To protect the patient against harmful germs, including the patient's own, from entering his/her body
3	AFTER BODY FLUID EXPOSURE RISK	WHEN? Clean your hands immediately after an exposure risk to body fluids (and after glove removal) WHY? To protect yourself and the healthcare environment from harmful patient germs
4	AFTER PATIENT CONTACT	WHEN? Clean your hands after touching a patient and her/his immediate surroundings when leaving the patient's side WHY? To protect yourself and the healthcare environment from harmful patient germs
5	AFTER CONTACT WITH PATIENT SURROUNDINGS	WHEN? Clean your hands after touching any object or furniture in the patient's immediate surroundings when leaving - even if the patient has not been touched WHY? To protect yourself and the healthcare environment from harmful patient germs

Adapted from WHO World Alliance for Patient Safety 2006



10. EQUALITY & DIVERSITY IMPACT ASSESSMENT

In reviewing this policy, the HR Policy Group considered, as a minimum, the following questions:

- Are the aims of this policy clear?
- Are responsibilities clearly identified?
- Has the policy been reviewed to ascertain any potential discrimination?
- Are there any specific groups impacted upon?
- Is this impact positive or negative?
- Could any impact constitute unlawful discrimination?
- Are communication proposals adequate?
- Does training need to be given? If so is this planned?

Adverse impact has been considered for age, disability, gender reassignment, marriage and civil partnership, pregnancy and maternity, race, religion and belief, sex, sexual orientation.

Blank version of the full Equality & Diversity Impact assessment can be found here:

http://eccho/Home/FormsGuidance.aspx?udt_575_param_index=E&udt_575_param_page=2

11. DOCUMENT CONTROL

Version Date	Version No.	Author/ Reviewer	Comments
July 2008	5	IPC	
June 2010	6	IPC	Change HH Technique from 6 step to 7. Revised HH Audit tool
May 2012	7	IPC	Revised HH Audit Tool Update References
March 2014	8	IPC	Revised HH Audit Tool
March 2016	9	IPC	Minor changes
March 2018	10	IPC	Minor changes
March 2020	11	IPC	Minor changes
June 2022	12	IPC	Preferred use of ABHR in clinical settings with stated exceptions

DOCUMENT CONTROL SHEET

Name of Document:	Hand Hygiene Policy
Version:	12
File Location / Document Name:	ECCHO
Date Of This Version:	June 2022
Produced By (Designation):	Infection Prevention and Control Team

Reviewed By:	(appropriate groups/virtually etc)
Synopsis And Outcomes Of Consultation Undertaken:	Changes relating to relevant committees/groups involved in ratification processes.
Synopsis And Outcomes Of Equality and Diversity Impact Assessment:	No specific issues
Ratified By (Committee):-	IPACC
Date Ratified:	June 2022
Distribute To:	All ECCH staff and contractors
Date Due For Review:	July 2024
Enquiries To:	infectionprevention@ecchcic.nhs.uk
Approved by Appropriate Group/Committee Approved by Policy Group Presented to IGC for information	<input type="checkbox"/> Date: <input type="checkbox"/> Date: <input type="checkbox"/> Date: